



Existence and Optimality of Cournot-Nash Equilibria in a Bilateral Oligopoly with Atoms and an Atomless Part

Francesca Busetto, Giulio Codognato,
Sayantan Ghosal, Ludovic Julien, Simone Tonin

November 2017

n. 6 / 2017

Existence and Optimality of Cournot-Nash Equilibria in a Bilateral Oligopoly with Atoms and an Atomless Part

Francesca Busetto*, Giulio Codognato†, Sayantan Ghosal‡
Ludovic Julien§, Simone Tonin¶

November 2017

Abstract

We consider a bilateral oligopoly version of the Shapley window model with large traders, represented as atoms, and small traders, represented by an atomless part. We first show the existence of a Cournot-Nash equilibrium. In order to cover the case where one of the two commodities is held only by atoms, our proof combines the existence proof provided by Busetto et al. (2017) for a mixed version of the Shapley window model with any finite number of commodities, each held by the atomless part, with the existence proof of a Cournot-Nash equilibrium for a strategic market game with a finite number of traders provided by Dubey and Shubik (1978). We then show, using a corollary proved by Shitovitz (1973), that a Cournot-Nash allocation is Pareto optimal if and only if it is a Walras allocation.

Journal of Economic Literature Classification Numbers: C72, D51.

*Dipartimento di Scienze Economiche e Statistiche, Università degli Studi di Udine, 33100 Udine, Italy.

†Dipartimento di Scienze Economiche e Statistiche, Università degli Studi di Udine, 33100 Udine, Italy, and Economix, UPL, Univ Paris Nanterre, CNRS, F92000 Nanterre, France.

‡Adam Smith Business School, University of Glasgow, Glasgow G12 8QQ, United Kingdom.

§Economix, UPL, Univ Paris Nanterre, CNRS, F92000 Nanterre, France.

¶Durham Business School, Durham University, Durham DH1 3LB, United Kingdom.

1 Introduction

Gabszewicz and Michel (1997) introduced the so-called model of bilateral oligopoly, which consists of a two-commodity exchange economy where each trader holds only one of the two commodities, whose aggregate amount is strictly positive in the economy. In this framework, traders' strategic interaction was modeled as in strategic market games à la Shapley and Shubik (see Giraud (2003) for a survey of this literature). This model was analyzed, in the case of a finite number of traders, by Bloch and Ghosal (1997), Bloch and Ferrer (2001), Dickson and Hartley (2008), Amir and Bloch (2009), among others.

In this paper, we consider the mixed bilateral oligopoly model introduced by Codognato et al. (2015). In this model, following Shitovitz (1973), there are large traders, represented as atoms, and small traders, represented by an atomless part. Moreover, traders' strategic interaction is formalized as in the Shapley window model, a strategic market game which was first proposed informally by Lloyds S. Shapley and further analyzed by Sahi and Yao (1989), Codognato and Ghosal (2000), Busetto et al. (2011), Busetto et al. (2017), among others.

The first goal of the paper is to prove the existence of a Cournot-Nash equilibrium for the mixed bilateral oligopoly version of the Shapley window model. Busetto et al. (2011) provided an existence proof for the mixed version of the Shapley window model with any finite number of commodities. Their proof is based on the same assumptions as the proof provided by Sahi and Yao (1989) in the case of exchange economies with a finite number of traders. In particular, it requires that there are at least two atoms with strictly positive endowments, continuously differentiable utility functions, and indifference curves contained in the strict interior of the commodity space: these restrictions are stated by Busetto et al. (2011) in their Assumption 4. Clearly, this proof does not apply to the bilateral oligopoly case where all atoms hold only one of the two commodities. Busetto et al. (2017) proposed an alternative existence proof which is essentially based on restrictions on endowments and preference of the atomless part of the economy rather than on atoms. In particular, they kept all the assumptions made by Busetto et al. (2011) with the exception of their Assumption 4 which was replaced by a new one requiring that the set of commodities is strongly connected through the characteristics of traders belonging to the atomless part. The existence proof in Busetto et al. (2017) used a theorem which shows that any sequence of prices corresponding to a sequence

of Cournot-Nash equilibria has a subsequence which converges to a strictly positive price vector and it requires that each commodity is held by a subset of the atomless part with positive measure. An appealing feature of our framework is the case where a commodity is held only by atoms. Therefore, in order to cover this case, we cannot directly use the price convergence theorem shown by Busetto et al. (2017) but we have to combine that proof with the price convergence result proved by Dubey and Shubik (1978) which holds for a strategic market game with a finite number of traders, i.e., for a purely atomic exchange economy. This is one reason why our existence proof is not just a two-commodity case of that provided by Busetto et al. (2017). The other reason is that we do not need that both of the two ordered pairs generated by the two commodities are connected through traders' characteristics to guarantee that the aggregate matrix of the bids obtained as the limit of a sequence of perturbed Cournot-Nash equilibria is irreducible, as in Busetto et al. (2017): only one of the two pairs must be connected. Therefore, Assumption 4 in Busetto et al. (2017) is more restrictive than its reformulation we use in this paper.

In the second main theorem of the paper, we provide a characterization of the Pareto optimality of Cournot-Nash allocations. The issue of Pareto optimality in strategic market games was raised since the seminal paper by Shapley and Shubik (1977). This first analysis was mainly an intuitive discussion of Pareto optimality in the Edgeworth box. Then, more formal results on this issue were obtained by Dubey (1980), Dubey et al. (1980), Aghion (1985), Dubey and Rogawski (1990), among others. These results were obtained using the approach to general equilibrium based on differential topology and hold generically. Our second theorem is a general result which holds under the same assumptions of the existence theorem with the only further restriction that both pairs of commodities are connected through the characteristics of the same subset of traders. It states that a Cournot-Nash allocation is Pareto optimal if and only if it is a Walras allocation, thereby establishing a relationship among the Cournotian tradition of oligopoly, the Walrasian tradition of perfect competition, and the Paretian analysis of optimality. Some examples provided by Codognato et al. (2015) show that this characterization holds non-vacuously.

The paper is organized as follows. In Section 2, we introduce the mathematical model. In section 3, we prove the existence of a Cournot-Nash equilibrium. In Section 4, we characterize the Pareto optimality of Cournot-Nash equilibria. In Section 5, we discuss the model. In Section 6, we draw some conclusions and we sketch some further lines of research.

2 Mathematical model

We consider a pure exchange economy with large traders, represented as atoms, and small traders, represented by an atomless part. The space of traders is denoted by the measure space (T, \mathcal{T}, μ) , where T is the set of traders, \mathcal{T} is the σ -algebra of all μ -measurable subsets of T , and μ is a real valued, non-negative, countably additive measure defined on \mathcal{T} . We assume that (T, \mathcal{T}, μ) is finite, i.e., $\mu(T) < \infty$. This implies that the measure space (T, \mathcal{T}, μ) contains at most countably many atoms. Let T_1 denote the set of atoms and T_0 the atomless part of T . We assume that $\mu(T_1) > 0$ and $\mu(T_0) > 0$. A null set of traders is a set of measure 0. Null sets of traders are systematically ignored throughout the paper. Thus, a statement asserted for “each” trader in a certain set is to be understood to hold for all such traders except possibly for a null set of traders. A coalition is a nonnull element of \mathcal{T} . The word “integrable” is to be understood in the sense of Lebesgue.

In the exchange economy, there are 2 different commodities. A commodity bundle is a point in R_+^2 . An assignment (of commodity bundles to traders) is an integrable function $\mathbf{x}: T \rightarrow R_+^2$. There is a fixed initial assignment \mathbf{w} , satisfying the following assumption.

Assumption 1. *There is a coalition S such that $\mu(S \cap T) > 0$, $\mu(S^c \cap T) > 0$, $\mathbf{w}^1(t) > 0$, $\mathbf{w}^2(t) = 0$, for each $t \in S$, $\mathbf{w}^1(t) = 0$, $\mathbf{w}^2(t) > 0$, for each $t \in S^c$. Moreover, $\text{card}(S \cap T_1) \geq 2$, whenever $\mu(S \cap T_0) = 0$, and $\text{card}(S^c \cap T_1) \geq 2$, whenever $\mu(S^c \cap T_0) = 0$.¹*

An allocation is an assignment \mathbf{x} for which $\int_T \mathbf{x}(t) d\mu = \int_T \mathbf{w}(t) d\mu$. The preferences of each trader $t \in T$ are described by a utility function $u_t: R_+^2 \rightarrow R$, satisfying the following assumptions.

Assumption 2. *$u_t: R_+^2 \rightarrow R$ is continuous, strongly monotone, and quasi-concave, for each $t \in T$.*

Let \mathcal{B} denote the Borel σ -algebra of R_+^2 . Moreover, let $\mathcal{T} \otimes \mathcal{B}$ denote the σ -algebra generated by the sets $E \times F$ such that $E \in \mathcal{T}$ and $F \in \mathcal{B}$.

Assumption 3. *$u: T \times R_+^2 \rightarrow R$, given by $u(t, x) = u_t(x)$, for each $t \in T$ and for each $x \in R_+^2$, is $\mathcal{T} \otimes \mathcal{B}$ -measurable.*

In order to state our last assumption, we need a preliminary definition. We say that commodities i, j stand in relation Q if there is a coalition T^i , such that $T^i = \{t \in T_0 : \mathbf{w}^i(t) > 0, \mathbf{w}^j(t) = 0\}$, $u_t(\cdot)$ is differentiable,

¹ $\text{card}(A)$ denotes the cardinality of a set A .

additively separable, i.e., $u_t(x) = v_t^i(x^i) + v_t^j(x^j)$, for each $x \in R_+^2$, and $\frac{dv_t^j(0)}{dx^j} = +\infty$, for each $t \in T^i$.² We can now introduce the last assumption.

Assumption 4. *Commodities 1 and 2 or commodities 2 and 1 stand in relation Q.*

A price vector is a nonnull vector $p \in R_+^2$. We say that a price vector p is normalized when $p \in \Delta$ where $\Delta = \{p \in R_+^2 : \sum_{i=1}^2 p^i = 1\}$.

Let $\mathbf{X}^0 : T_0 \times R_{++}^2 \rightarrow \mathcal{P}(R_+^2)$ be a correspondence such that, for each $t \in T_0$ and for each $p \in R_{++}^2$, $\mathbf{X}^0(t, p) = \operatorname{argmax}\{u(x) : x \in R_+^2 \text{ and } px \leq p\mathbf{w}(t)\}$. For each $p \in R_{++}^2$, let $\int_{T_0} \mathbf{X}^0(t, p) d\mu = \{\int_{T_0} \mathbf{x}^0(t, p) d\mu : \mathbf{x}^0(\cdot, p) \text{ is integrable and } \mathbf{x}^0(t, p) \in \mathbf{X}^0(t, p), \text{ for each } t \in T_0\}$. Finally, let $\mathbf{Z}^0 : R_{++}^2 \rightarrow \mathcal{P}(R^2)$ be a correspondence which associates with each $p \in R_{++}^2$ the Minkowski difference between the set $\int_{T_0} \mathbf{X}^0(t, p) d\mu$ and the set $\{\int_{T_0} \mathbf{w}(t) d\mu\}$.³ According to Debreu (1982), let $|x| = \sum_{i=1}^2 |x^i|$, for each $x \in R^2$, and let $d[0, V] = \inf_{x \in V} |x|$, for each $V \subset R^2$. The next proposition, which we shall use in the proof of the existence theorem, is based on Property (iv) in Debreu (1982) p. 728.

Proposition 1. *Let $\{p^n\}$ be a sequence of normalized price vectors such that $p^n \gg 0$, for each $n = 1, 2, \dots$, which converges to a normalized price vector \bar{p} . Then, $\bar{p}^1 = 0$ and $\mu(S^c \cap T_0) > 0$, or, $\bar{p}^2 = 0$ and $\mu(S \cap T_0) > 0$, imply that the sequence $\{d[0, \mathbf{Z}^0(p^n)]\}$ diverges to $+\infty$.*

Proof. Let $\{p^n\}$ be a sequence of normalized price vectors such that $p^n \gg 0$, for each $n = 1, 2, \dots$, which converges to a normalized price vector \bar{p} . Suppose that $\bar{p}^1 = 0$ and $\mu(S^c \cap T_0) > 0$. Then, we have that $\bar{p}^2 = 1$. But then, the sequence $\{d[0, \mathbf{X}^0(t, p^n)]\}$ diverges to $+\infty$ as $\bar{p}^2 \mathbf{w}^2(t) > 0$, for each $t \in S^c \cap T_0$, by Lemma 4 in Debreu (1982) p. 721. Therefore, $\{d[0, \mathbf{Z}^0(p^n)]\}$ diverges to $+\infty$ as $\mu(S^c \cap T_0) > 0$, by the argument used in the proof of Property (iv) in Debreu (1982) p. 728. Suppose that $\bar{p}^2 = 0$ and $\mu(S \cap T_0) > 0$. Then, $\{d[0, \mathbf{Z}^0(p^n)]\}$ diverges to $+\infty$, by using symmetrically the previous argument. Hence, $\bar{p}^1 = 0$ and $\mu(S^c \cap T_0) > 0$, or, $\bar{p}^2 = 0$ and $\mu(S \cap T_0) > 0$, imply that the sequence $\{d[0, \mathbf{Z}^0(p^n)]\}$ diverges to $+\infty$. ■

A Walras equilibrium is a pair (p, \mathbf{x}) , consisting of a price vector p and an allocation \mathbf{x} , such that $px(t) = p\mathbf{w}(t)$ and $u_t(\mathbf{x}(t)) \geq u_t(y)$, for all $y \in$

²In this definition, differentiability means continuous differentiability and it should be understood to include the case of infinite partial derivatives along the boundary of the consumption set (for a discussion of this case, see, for instance, Kreps (2012), p. 58).

³For a discussion of the properties of the correspondences introduced above and their proofs see, for instance, Debreu (1982), Section 4.

$\{x \in R_+^2 : px = p\mathbf{w}(t)\}$, for each $t \in T$. A Walras allocation is an allocation \mathbf{x} for which there exists a price vector p such that the pair (p, \mathbf{x}) is a Walras equilibrium.

We now introduce the two-commodity version of the Shapley window model proposed by Codognato et al. (2015). A strategy correspondence is a correspondence $\mathbf{B} : T \rightarrow \mathcal{P}(R_+^4)$ such that, for each $t \in T$, $\mathbf{B}(t) = \{(b_{ij}) \in R_+^4 : \sum_{j=1}^2 b_{ij} \leq \mathbf{w}^i(t), i = 1, 2\}$. With some abuse of notation, we denote by $b(t) \in \mathbf{B}(t)$ a strategy of trader t , where $b_{ij}(t)$, $i, j = 1, 2$, represents the amount of commodity i that trader t offers in exchange for commodity j . A strategy selection is an integrable function $\mathbf{b} : T \rightarrow R_+^4$, such that, for each $t \in T$, $\mathbf{b}(t) \in \mathbf{B}(t)$. Given a strategy selection \mathbf{b} , we call aggregate matrix the matrix $\bar{\mathbf{B}}$ such that $\bar{b}_{ij} = (\int_T \mathbf{b}_{ij}(t) d\mu)$, $i, j = 1, 2$. Moreover, we denote by $\mathbf{b} \setminus b(t)$ the strategy selection obtained from \mathbf{b} by replacing $\mathbf{b}(t)$ with $b(t) \in \mathbf{B}(t)$ and by $\bar{\mathbf{B}} \setminus b(t)$ the corresponding aggregate matrix.

We then introduce three further definitions (see Sahi and Yao (1989)).

Definition 1. A nonnegative square matrix A is said to be irreducible if, for every pair (i, j) , with $i \neq j$, there is a positive integer k such that $a_{ij}^{(k)} > 0$, where $a_{ij}^{(k)}$ denotes the ij -th entry of the k -th power A^k of A .

Definition 2. A nonnegative square matrix A is said to be completely reducible if, after a permutation of indices, it can be written in a block-diagonal form such that each diagonal block is irreducible.

Definition 3. Given a strategy selection \mathbf{b} , a price vector p is said to be market clearing if

$$p \in R_{++}^2, \sum_{i=1}^2 p^i \bar{b}_{ij} = p^j (\sum_{i=1}^2 \bar{b}_{ji}), j = 1, 2. \quad (1)$$

By Lemma 1 in Sahi and Yao (1989), there is a unique, up to a scalar multiple, price vector p satisfying (1) if and only if $\bar{\mathbf{B}}$ is irreducible. Then, we denote by $p(\mathbf{b})$ a function which associates with each strategy selection \mathbf{b} the unique, up to a scalar multiple, price vector p satisfying (1), if $\bar{\mathbf{B}}$ is irreducible, and is equal to 0, otherwise. For each strategy selection \mathbf{b} such that $p(\mathbf{b}) \gg 0$, we assume that the price vector $p(\mathbf{b})$ is normalized.

Given a strategy selection \mathbf{b} and a price vector p , consider the assignment determined as follows:

$$\mathbf{x}^j(t, \mathbf{b}(t), p) = \mathbf{w}^j(t) - \sum_{i=1}^2 \mathbf{b}_{ji}(t) + \sum_{i=1}^2 \mathbf{b}_{ij}(t) \frac{p^i}{p^j}, \text{ if } p \in R_{++}^2,$$

$$\mathbf{x}^j(t, \mathbf{b}(t), p) = \mathbf{w}^j(t), \text{ otherwise,}$$

$j = 1, 2$, for each $t \in T$.

Given a strategy selection \mathbf{b} and the function $p(\mathbf{b})$, the traders' final holdings are determined according to this rule and consequently expressed by the assignment

$$\mathbf{x}(t) = \mathbf{x}(t, \mathbf{b}(t), p(\mathbf{b})),$$

for each $t \in T$.⁴ It is straightforward to show that this assignment is an allocation.

We are now able to define a notion of Cournot-Nash equilibrium for this reformulation of the Shapley window model (see Codognato and Ghosal (2000) and Busetto et al. (2011)).

Definition 4. *A strategy selection $\hat{\mathbf{b}}$ such that $\bar{\mathbf{B}}$ is irreducible is a Cournot-Nash equilibrium if*

$$u_t(\mathbf{x}(t, \hat{\mathbf{b}}(t), p(\hat{\mathbf{b}}))) \geq u_t(\mathbf{x}(t, b(t), p(\hat{\mathbf{b}} \setminus b(t)))),$$

for each $b(t) \in \mathbf{B}(t)$ and for each $t \in T$.

A Cournot-Nash allocation is an allocation $\hat{\mathbf{x}}$ such that $\hat{\mathbf{x}}(t) = \mathbf{x}(t, \hat{\mathbf{b}}(t), p(\hat{\mathbf{b}}))$, for each $t \in T$, where $\hat{\mathbf{b}}$ is a Cournot-Nash equilibrium.

Finally, we introduce a perturbation of the strategic market game which was used by Sahi and Yao (1989) and Busetto et al. (2011) for their existence proofs. Given $\epsilon > 0$ and a strategy selection \mathbf{b} , we define the aggregate matrix $\bar{\mathbf{B}}_\epsilon$ to be the matrix such that $\bar{\mathbf{b}}_{\epsilon ij} = (\bar{\mathbf{b}}_{ij} + \epsilon)$, $i, j = 1, 2$. Clearly, the matrix $\bar{\mathbf{B}}_\epsilon$ is irreducible. The interpretation is that an outside agency places fixed bids of ϵ for each pair of commodities 1 and 2. Given $\epsilon > 0$, we denote by $p^\epsilon(\mathbf{b})$ the function which associates, with each strategy selection \mathbf{b} , the unique, up to a scalar multiple, price vector which satisfies

$$\sum_{i=1}^2 p^i(\bar{\mathbf{b}}_{ij} + \epsilon) = p^j(\sum_{i=1}^2 (\bar{\mathbf{b}}_{ji} + \epsilon)), \quad j = 1, 2. \quad (2)$$

For each strategy selection \mathbf{b} , we assume that the price vector $p^\epsilon(\mathbf{b})$ is normalized.

⁴In order to save in notation, with some abuse, we denote by \mathbf{x} both the function $\mathbf{x}(t)$ and the function $\mathbf{x}(t, \mathbf{b}(t), p(\mathbf{b}))$.

Definition 5. Given $\epsilon > 0$, a strategy selection $\hat{\mathbf{b}}^\epsilon$ is an ϵ -Cournot-Nash equilibrium if

$$u_t(\mathbf{x}(t, \hat{\mathbf{b}}^\epsilon(t), p^\epsilon(\hat{\mathbf{b}}^\epsilon))) \geq u_t(\mathbf{x}(t, b(t), p^\epsilon(\hat{\mathbf{b}}^\epsilon \setminus b(t)))),$$

for each $b(t) \in \mathbf{B}(t)$ and for each $t \in T$.

3 Existence

In this section we provide a proof of the existence of a Cournot-Nash equilibrium, under the assumptions listed above.

Theorem 1. Under Assumptions 1, 2, 3, and 4, there exists a Cournot-Nash equilibrium $\hat{\mathbf{b}}$.

Proof. To show Theorem 1, we first need to prove the existence of an ϵ -Cournot-Nash equilibrium. The following lemma, which was proved by Busetto et al. (2011) applying the Kakutani-Fan-Glicksberg theorem, states that such an equilibrium exists.

Lemma 1. For each $\epsilon > 0$, there exists an ϵ -Cournot-Nash equilibrium $\hat{\mathbf{b}}^\epsilon$.

Proof. See the proof of Lemma 3 in Busetto et al. (2011). ■

We now show that there exists the limit of a sequence of ϵ -Cournot-Nash equilibria and that this limit is a Cournot-Nash equilibrium. Let $\epsilon_n = \frac{1}{n}$, $n = 1, 2, \dots$. By Lemma 1, for each $n = 1, 2, \dots$, there is an ϵ -Cournot-Nash equilibrium $\hat{\mathbf{b}}^{\epsilon_n}$. The next step would consist in showing that any sequence of normalized prices generated by the sequence of ϵ -Cournot-Nash equilibria corresponding to the sequence $\{\epsilon_n\}$ has a convergent subsequence whose limit is a strictly positive normalized price vector. In order to prove this result, we cannot use the price convergence theorem proved by Busetto et al. (2017) as the proof of this theorem requires that each commodity is held by a subset of the atomless part with positive measure whereas, in our framework, a commodity may be held only by atoms. Therefore, we need to combine the proof of the price convergence theorem provided by Busetto et al. (2017) with the price convergence result proved by Dubey and Shubik (1978) which holds for a purely atomic exchange economy. To this end, we need the following preliminary lemma which is based on the uniform monotonicity lemma proved by Dubey and Shubik (1978).

Lemma 2. Consider an atom $\tau \in T_1$ and a commodity $j \in \{1, 2\}$. For each real number $H > 0$, there is a real number $0 < h(u_\tau(\cdot), j, H) < 1$, depending on $u_\tau(\cdot)$, j , and H , such that if $x \in R_+^2$, $\|x\| \leq H$, $y \in R_+^2$ and $\|y - x\| \leq h(u_\tau(\cdot), j, H)$, then $u_\tau(y + e^j) > u_\tau(x)$.⁵

Proof. It is an immediate consequence of Lemma C (the uniform monotonicity lemma) in Dubey and Shubik (1978) as $u_\tau(\cdot)$ is continuous and strongly monotone, by Assumption 2. ■

We can now state and prove the price convergence lemma.

Lemma 3. Let $\{\hat{p}^{\epsilon_n}\}$ be a sequence of normalized prices such that $\{\hat{p}^{\epsilon_n}\} = p(\hat{\mathbf{b}}^{\epsilon_n})$ where $\hat{\mathbf{b}}^{\epsilon_n}$ is an ϵ -Cournot-Nash equilibrium, for each $n = 1, 2, \dots$. Then, there exists a subsequence $\{\hat{p}^{\epsilon_{k_n}}\}$ of the sequence $\{\hat{p}^{\epsilon_n}\}$ which converges to a normalized price vector $\hat{p} \gg 0$.

Proof. Let $\{\hat{p}^{\epsilon_n}\}$ be a sequence of normalized prices such that $\{\hat{p}^{\epsilon_n}\} = p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n})$ where $\hat{\mathbf{b}}^{\epsilon_n}$ is an ϵ -Cournot-Nash equilibrium, for each $n = 1, 2, \dots$. Then, there is a subsequence $\{\hat{p}^{\epsilon_{k_n}}\}$ of the sequence $\{\hat{p}^{\epsilon_n}\}$ which converges to a price vector $\hat{p} \in \Delta$ as the unit simplex Δ is a compact set. Consider the case where $\mu(S \cap T_0) > 0$ and $\mu(S^c \cap T_0) > 0$. Suppose that $\hat{p}^1 = 0$. Then, the sequence $\{d[0, \mathbf{Z}^0(\hat{p}^{\epsilon_{k_n}})]\}$ diverges to $+\infty$ as $\mu(S^c \cap T_0) > 0$, by Proposition 1. We now adapt the argument used by Busetto et al. (2017) to prove their Theorem 1 to our framework. Let $\hat{\mathbf{x}}^{\epsilon_n}(t) = \mathbf{x}(t, \hat{\mathbf{b}}^{\epsilon_n}(t), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n}))$, for each $t \in T$, and for each $n = 1, 2, \dots$. Then, $\hat{\mathbf{x}}^{\epsilon_n}(t) \in \mathbf{X}^0(t, p^{\epsilon_n})$, for each $t \in T_0$, and for each $n = 1, 2, \dots$, by the same argument used by Codognato and Ghosal (2000) to prove their Theorem 2. But then, $(\int_{T_0} \hat{\mathbf{x}}^{\epsilon_n}(t) d\mu - \int_{T_0} \mathbf{w}(t) d\mu) \in \mathbf{Z}^0(\hat{p}^{\epsilon_n})$, for each $n = 1, 2, \dots$. We have that

$$\int_{T_0} \hat{\mathbf{x}}^{\epsilon_n}(t) d\mu \leq \int_{T_0} \mathbf{w}(t) d\mu + \int_{T_1} \mathbf{w}(t) d\mu + e^1 + e^2$$

as $\int_T \hat{\mathbf{x}}^{\epsilon_n}(t) d\mu \leq \int_T \mathbf{w}(t) d\mu + \epsilon_n e^1 + \epsilon_n e^2$, for each $n = 1, 2, \dots$. Then,

$$\left| \int_{T_0} \hat{\mathbf{x}}^{i\epsilon_n}(t) d\mu - \int_{T_0} \mathbf{w}^i(t) d\mu \right| \leq \int_{T_0} \mathbf{w}^i(t) d\mu + \int_{T_1} \mathbf{w}^i(t) d\mu + 1$$

as $-\int_{T_1} \mathbf{w}^i(t) d\mu - 1 \leq \int_{T_0} \hat{\mathbf{x}}^{i\epsilon_n}(t) d\mu \leq 2 \int_{T_0} \mathbf{w}^i(t) d\mu + \int_{T_1} \mathbf{w}^i(t) d\mu + 1$, $i = 1, 2$, for each $n = 1, 2, \dots$. But then,

$$\sum_{i=1}^2 \left| \int_{T_0} \hat{\mathbf{x}}^{i\epsilon_n}(t) d\mu - \int_{T_0} \mathbf{w}^i(t) d\mu \right| \leq \sum_{i=1}^2 \left(\int_{T_0} \mathbf{w}^i(t) d\mu + \int_{T_1} \mathbf{w}^i(t) d\mu + 1 \right),$$

⁵ $\|\cdot\|$ denotes the Euclidean norm and e^j denotes the vector in R^2 whose j th coordinate is 1 and whose other coordinate vanishes.

for each $n = 1, 2, \dots$. Moreover, there exists an n_0 such that

$$d[0, \mathbf{Z}^0(\hat{p}^{\epsilon_{k_n}})] > \sum_{i=1}^2 \left(\int_{T_0} \mathbf{w}^i(t) d\mu + \int_{T_1} \mathbf{w}^i(t) d\mu + 1 \right),$$

for each $n \geq n_0$, as the sequence $\{d[0, \mathbf{Z}^0(\hat{p}^{\epsilon_{k_n}})]\}$ diverges to $+\infty$. Then,

$$\sum_{i=1}^2 \left| \int_{T_0} \hat{\mathbf{x}}^{i\epsilon_{k_n}}(t) d\mu - \int_{T_0} \mathbf{w}^i(t) d\mu \right| > \sum_{i=1}^2 \left(\int_{T_0} \mathbf{w}^i(t) d\mu + \int_{T_1} \mathbf{w}^i(t) d\mu + 1 \right)$$

as $\sum_{i=1}^2 \left| \int_{T_0} \hat{\mathbf{x}}^{i\epsilon_{k_n}}(t) d\mu - \int_{T_0} \mathbf{w}^i(t) d\mu \right| \geq d[0, \mathbf{Z}^0(\hat{p}^{\epsilon_{k_n}})]$, for each $n \geq n_0$, a contradiction. Therefore, we must have that $\hat{p}^1 > 0$. Suppose that $\hat{p}^2 = 0$. Then, by using symmetrically the previous argument, we generate the same contradiction. Therefore, we must have that $\hat{p}^2 > 0$. Consider the case where $\mu(S \cap T_0) = 0$ or $\mu(S^c \cap T_0) = 0$. Suppose, without loss of generality, that $\mu(S \cap T_0) = 0$. Then, we have that $\mu(S^c \cap T_0) > 0$ as $\mu(T_0) > 0$. Moreover, there are at least two atoms $\tau, \rho \in S \cap T_1$, by Assumption 1. We have that $\hat{p}^1 > 0$, as $\mu(S^c \cap T_0) > 0$, by the same argument used in the proof of the previous case. In order to prove that $\hat{p}^2 > 0$, we now show that there is a real number $\eta > 0$ such that

$$\frac{\hat{p}^{2\epsilon_n}}{\hat{p}^{1\epsilon_n}} > \eta, \quad (3)$$

for each $n = 1, 2, \dots$. To this end, we adapt to our framework the proof of Lemma 2 provided by Dubey and Shubik (1978). In what follows, we shall use the fact that (2) and the normalization rule imply straightforwardly that (3) holds if and only if

$$\frac{\bar{\mathbf{b}}_{12}^{\epsilon_n} + \epsilon_n}{\bar{\mathbf{b}}_{21}^{\epsilon_n} + \epsilon_n} > \eta,$$

for each $n = 1, 2, \dots$. Consider any n . We now prove that $\hat{\mathbf{b}}_{12}^{\epsilon_n}(\tau) \leq \frac{\bar{\mathbf{b}}_{12}^{\epsilon_n}}{2}$ or $\hat{\mathbf{b}}_{12}^{\epsilon_n}(\rho) \leq \frac{\bar{\mathbf{b}}_{12}^{\epsilon_n}}{2}$. Suppose, by way of contradiction, that $\hat{\mathbf{b}}_{12}^{\epsilon_n}(\tau) > \frac{\bar{\mathbf{b}}_{12}^{\epsilon_n}}{2}$ and $\hat{\mathbf{b}}_{12}^{\epsilon_n}(\rho) > \frac{\bar{\mathbf{b}}_{12}^{\epsilon_n}}{2}$. Then, we have that $\hat{\mathbf{b}}_{12}^{\epsilon_n}(\tau) + \hat{\mathbf{b}}_{12}^{\epsilon_n}(\rho) > \bar{\mathbf{b}}_{12}^{\epsilon_n}$, a contradiction. Therefore, we must have that $\hat{\mathbf{b}}_{12}^{\epsilon_n}(\tau) \leq \frac{\bar{\mathbf{b}}_{12}^{\epsilon_n}}{2}$ or $\hat{\mathbf{b}}_{12}^{\epsilon_n}(\rho) \leq \frac{\bar{\mathbf{b}}_{12}^{\epsilon_n}}{2}$. Suppose that $\hat{\mathbf{b}}_{12}^{\epsilon_n}(\tau) \leq \frac{\bar{\mathbf{b}}_{12}^{\epsilon_n}}{2}$. Moreover, suppose that $\mathbf{w}^1(\tau) - \hat{\mathbf{b}}_{12}^{\epsilon_n}(\tau) \geq \frac{\mathbf{w}^1(\tau)}{2}$. Let

$0 < \gamma < \min\{\epsilon_n, \frac{\mathbf{w}^1(\tau)}{2}, 2\frac{\bar{\mathbf{b}}_{12}^{\epsilon_n} + \epsilon_n}{\bar{\mathbf{b}}_{21}^{\epsilon_n} + \epsilon_n}\}$ and let $b^\gamma(\tau) = \hat{\mathbf{b}}^{\epsilon_n}(\tau) + \gamma e^2$. Then, we have

$$\begin{aligned} & \mathbf{x}^1(\tau, b^\gamma(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n} \setminus b^\gamma(\tau))) - \mathbf{x}^1(\tau, \hat{\mathbf{b}}^{\epsilon_n}(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n})) \\ &= (\mathbf{w}^1 - \hat{\mathbf{b}}_{12}^{\epsilon_n} - \gamma) - (\mathbf{w}^1 - \hat{\mathbf{b}}_{12}^{\epsilon_n}) \\ &= -\gamma \end{aligned}$$

and

$$\begin{aligned} & \mathbf{x}^2(\tau, b^\gamma(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n} \setminus b^\gamma(\tau))) - \mathbf{x}^2(\tau, \hat{\mathbf{b}}^{\epsilon_n}(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n})) \\ &= (\hat{\mathbf{b}}_{12}^{\epsilon_n}(\tau) + \gamma) \frac{\bar{\mathbf{b}}_{21}^{\epsilon_n} + \epsilon_n}{\bar{\mathbf{b}}_{12}^{\epsilon_n} + \epsilon_n + \gamma} - \hat{\mathbf{b}}_{12}^{\epsilon_n}(\tau) \frac{\bar{\mathbf{b}}_{21}^{\epsilon_n} + \epsilon_n}{\bar{\mathbf{b}}_{12}^{\epsilon_n} + \epsilon_n} \\ &= \frac{\bar{\mathbf{b}}_{12}^{\epsilon_n} + \epsilon_n - \hat{\mathbf{b}}_{12}^{\epsilon_n}(\tau) \frac{\bar{\mathbf{b}}_{21}^{\epsilon_n} + \epsilon_n}{\bar{\mathbf{b}}_{12}^{\epsilon_n} + \epsilon_n}}{\bar{\mathbf{b}}_{12}^{\epsilon_n} + \epsilon_n + \gamma} \gamma \\ &> \frac{\frac{\bar{\mathbf{b}}_{12}^{\epsilon_n}}{2} + \frac{\epsilon_n}{2} + \frac{\gamma}{2} \frac{\bar{\mathbf{b}}_{21}^{\epsilon_n} + \epsilon_n}{\bar{\mathbf{b}}_{12}^{\epsilon_n} + \epsilon_n}}{\bar{\mathbf{b}}_{12}^{\epsilon_n} + \epsilon_n + \gamma} \gamma, \end{aligned}$$

as $\hat{\mathbf{b}}_{12}^{\epsilon_n}(\tau) \leq \frac{\bar{\mathbf{b}}_{12}^{\epsilon_n}}{2}$ and $\gamma < \epsilon_n$. Then, we obtain

$$\mathbf{x}^2(\tau, b^\gamma(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n} \setminus b^\gamma(\tau))) - \mathbf{x}^2(\tau, \hat{\mathbf{b}}^{\epsilon_n}(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n})) > \frac{1}{2} \frac{\bar{\mathbf{b}}_{21}^{\epsilon_n} + \epsilon_n}{\bar{\mathbf{b}}_{12}^{\epsilon_n} + \epsilon_n} \gamma. \quad (4)$$

Let us define

$$z = -2 \frac{\bar{\mathbf{b}}_{12}^{\epsilon_n} + \epsilon_n}{\bar{\mathbf{b}}_{21}^{\epsilon_n} + \epsilon_n} e^1.$$

Then, we have the vector inequality

$$\begin{aligned} & \mathbf{x}(\tau, b^\gamma(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n} \setminus b^\gamma(\tau))) \\ & \geq \mathbf{x}(\tau, \hat{\mathbf{b}}^{\epsilon_n}(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n})) + \frac{1}{2} \frac{\bar{\mathbf{b}}_{21}^{\epsilon_n} + \epsilon_n}{\bar{\mathbf{b}}_{12}^{\epsilon_n} + \epsilon_n} \gamma (z + e^2), \end{aligned} \quad (5)$$

where the inequality is strict for the second component by (4). Let $H = \sqrt{2} \max\{\int_T \mathbf{w}^1(t) d\mu + 1, \int_T \mathbf{w}^2(t) d\mu + 1\}$ and let $y = \mathbf{x}(\tau, \hat{\mathbf{b}}^{\epsilon_n}(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n})) + z$. It is straightforward to verify that $\mathbf{x}(\tau, \hat{\mathbf{b}}^{\epsilon_n}(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n})) \in R_+^2$ and

$\|\mathbf{x}(\tau, \hat{\mathbf{b}}^{\epsilon_n}(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n}))\| \leq H$. Suppose that $y \in R_+^2$ and $\|z\| \leq h(u_\tau(\cdot), 2, H)$. Then, by Lemma 2, we obtain that

$$u_\tau(\mathbf{x}(\tau, \hat{\mathbf{b}}^{\epsilon_n}(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n})) + z + e^2) > u_\tau(\mathbf{x}(\tau, \hat{\mathbf{b}}^{\epsilon_n}(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n}))).$$

But then, we have that

$$u_\tau(\mathbf{x}(\tau, \hat{\mathbf{b}}^{\epsilon_n}(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n})) + \frac{1}{2} \frac{\bar{\mathbf{b}}_{21}^{\epsilon_n} + \epsilon_n}{\bar{\mathbf{b}}_{21}^{\epsilon_n} + \epsilon_n} \gamma(z + e^2)) \geq u_\tau(\mathbf{x}(\tau, \hat{\mathbf{b}}^{\epsilon_n}(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n}))),$$

as $0 < \frac{1}{2} \frac{\bar{\mathbf{b}}_{21}^{\epsilon_n} + \epsilon_n}{\bar{\mathbf{b}}_{21}^{\epsilon_n} + \epsilon_n} \gamma < 1$ and the function $u_\tau(\cdot)$ is quasi-concave, by Assumption 2. Therefore, it follows that

$$u_\tau(\mathbf{x}(\tau, b^\gamma(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n} \setminus b^\gamma(\tau)))) > u_\tau(\mathbf{x}(\tau, \hat{\mathbf{b}}^{\epsilon_n}(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n}))),$$

as (5) holds strictly for its second component and $u_\tau(\cdot)$ is strongly monotone, by Assumption 2, a contradiction. Thus, it must be that $y \notin R_+^2$ or $\|z\| > h(u_\tau(\cdot), 2, H)$. Suppose that $y \notin R_+^2$. Then, it follows that

$$\mathbf{x}^1(\tau, \hat{\mathbf{b}}^{\epsilon_n}(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n})) - 2 \frac{\bar{\mathbf{b}}_{12}^{\epsilon_n} + \epsilon_n}{\bar{\mathbf{b}}_{21}^{\epsilon_n} + \epsilon_n} < 0,$$

as $y = \mathbf{x}(\tau, \hat{\mathbf{b}}^{\epsilon_n}(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n})) + z$ and $\mathbf{x}(\tau, \hat{\mathbf{b}}^{\epsilon_n}(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n})) \in R_+^2$. But then, it must be that

$$\frac{\bar{\mathbf{b}}_{12}^{\epsilon_n} + \epsilon_n}{\bar{\mathbf{b}}_{21}^{\epsilon_n} + \epsilon_n} > \frac{\mathbf{w}^1(\tau)}{4},$$

as $\mathbf{x}^1(\tau, \hat{\mathbf{b}}^{\epsilon_n}(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n})) = \mathbf{w}^1(\tau) - \hat{\mathbf{b}}_{12}^{\epsilon_n}(\tau) \geq \frac{\mathbf{w}^1(\tau)}{2}$. Suppose that $\|z\| > h(u_\tau(\cdot), 2, H)$. Then, we have that

$$\frac{\bar{\mathbf{b}}_{12}^{\epsilon_n} + \epsilon_n}{\bar{\mathbf{b}}_{21}^{\epsilon_n} + \epsilon_n} > \frac{h(u_\tau(\cdot), 2, H)}{2}.$$

Suppose now that $\mathbf{w}^1(\tau) - \hat{\mathbf{b}}_{12}^{\epsilon_n}(\tau) < \frac{\mathbf{w}^1(\tau)}{2}$. Then, we have that $\hat{\mathbf{b}}_{12}^{\epsilon_n}(\tau) > \frac{\mathbf{w}^1(\tau)}{2}$. But then, it must be that

$$\frac{\bar{\mathbf{b}}_{12}^{\epsilon_n} + \epsilon_n}{\bar{\mathbf{b}}_{21}^{\epsilon_n} + \epsilon_n} > \frac{\mathbf{w}^1(\tau)}{2(\int_T \mathbf{w}^2(t) d\mu + 1)}.$$

Let

$$\alpha = \min \left\{ \frac{\mathbf{w}^1(\tau)}{4}, \frac{h(u_\tau(\cdot), 2, H)}{2}, \frac{\mathbf{w}^1(\tau)}{2(\int_T \mathbf{w}^2(t) d\mu + 1)} \right\}.$$

Thus, we have that

$$\frac{\hat{p}^{2\epsilon_n}}{\hat{p}^{1\epsilon_n}} > \alpha.$$

Suppose that $\hat{\mathbf{b}}_{12}^{\epsilon_n}(\rho) \leq \frac{\bar{\mathbf{b}}_{12}^{\epsilon_n}}{2}$. Let

$$\beta = \min \left\{ \frac{\mathbf{w}^1(\rho)}{4}, \frac{h(u_\rho(\cdot), 2, H)}{2}, \frac{\mathbf{w}^1(\rho)}{2(\int_T \mathbf{w}^2(t) d\mu + 1)} \right\}.$$

Thus, by the same argument used in the previous case, we have that

$$\frac{\hat{p}^{2\epsilon_n}}{\hat{p}^{1\epsilon_n}} > \beta.$$

Let $\eta = \min\{\alpha, \beta\}$. Therefore, we can conclude that

$$\frac{\hat{p}^{2\epsilon_n}}{\hat{p}^{1\epsilon_n}} > \eta,$$

for each $n = 1, 2, \dots$. Consider the sequence $\{\hat{p}^{\epsilon_{k_n}}\}$. From (3), we obtain that

$$\hat{p}^{2\epsilon_{k_n}} > \eta \hat{p}^{1\epsilon_{k_n}},$$

for each $n = 1, 2, \dots$. Then, we obtain that

$$\hat{p}^2 > \eta \hat{p}^1,$$

as the sequence $\{\hat{p}^{\epsilon_{k_n}}\}$ converges to \hat{p} . But then, we have that $\hat{p}^2 > 0$ as $\eta > 0$ and $\hat{p}^1 > 0$. Hence, having considered all possible cases, we can conclude that $\hat{p} \gg 0$. \blacksquare

We now follow the argument used by Busetto et al. (2017) to prove their Theorem 2. In the next part of the proof, we apply a generalization of the Fatou lemma in several dimensions provided by Artstein (1979). By Lemma 1, there is an ϵ -Cournot-Nash equilibrium $\hat{\mathbf{b}}^{\epsilon_n}$, for each $n = 1, 2, \dots$. The fact that the sequence $\{\tilde{\mathbf{B}}^{\epsilon_n}\}$ belongs to the compact set $\{(b_{ij}) \in R_+^4 : b_{ij} \leq \int_T \mathbf{w}^i(t) d\mu, i, j = 1, 2\}$ and the sequence $\{\hat{p}^{\epsilon_n}\}$, where $\hat{p}^{\epsilon_n} = p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n})$, belongs to the unit simplex Δ , for each $n = 1, 2, \dots$, implies that there is a subsequence $\{\tilde{\mathbf{B}}^{\epsilon_{k_n}}\}$ of the sequence $\{\tilde{\mathbf{B}}^{\epsilon_n}\}$ which converges to an element

of the set $\{(b_{ij}) \in R_+^4 : b_{ij} \leq \int_T \mathbf{w}^i(t) d\mu, i, j = 1, 2\}$ and a subsequence $\{\hat{p}^{\epsilon_{k_n}}\}$ of the sequence $\{\hat{p}^{\epsilon_n}\}$ which converges to a price vector $\hat{p} \in \Delta$, with $\hat{p} \gg 0$, by Lemma 3. Since the sequence $\{\hat{\mathbf{b}}^{\epsilon_{k_n}}\}$ satisfies the assumptions of Theorem A in Artstein (1979), by this theorem there is a function $\hat{\mathbf{b}}$ such that $\hat{\mathbf{b}}(t)$ is a limit point of the sequence $\{\hat{\mathbf{b}}^{\epsilon_{k_n}}(t)\}$, for each $t \in T$, and such that the sequence $\{\tilde{\mathbf{B}}^{\epsilon_{k_n}}\}$ converges to $\tilde{\mathbf{B}}$. Moreover, \hat{p} and $\tilde{\mathbf{B}}$ satisfy (1) as $\hat{p}^{\epsilon_{k_n}}$ and $\tilde{\mathbf{B}}^{\epsilon_{k_n}}$ satisfy (2), for each $n = 1, 2, \dots$, the sequence $\{\hat{p}^{\epsilon_{k_n}}\}$ converges to \hat{p} , the sequence $\{\tilde{\mathbf{B}}^{\epsilon_{k_n}}\}$ converges to $\tilde{\mathbf{B}}$, and the sequence $\{\epsilon_{k_n}\}$ converges to 0. Then, the matrix $\tilde{\mathbf{B}}$ is completely reducible, by Lemma 1 in Sahi and Yao (1989), as $\hat{p} \gg 0$. We want now to prove that $\tilde{\mathbf{B}}$ must be irreducible. Suppose, without loss of generality, that commodities 2 and 1 stand in the relation Q . We now show that $\tilde{\mathbf{b}}_{21} > 0$. Suppose that $\tilde{\mathbf{b}}_{21} = 0$. Then, we have that $\int_{T_2} \hat{\mathbf{b}}_{21}(t) d\mu = 0$ as $\mu(T^2) > 0$. Consider a trader $\tau \in T^2$. We can suppose that $\hat{\mathbf{b}}_{21}(\tau) = 0$ as we ignore null sets. Since $\hat{\mathbf{b}}(\tau)$ is a limit point of the sequence $\{\hat{\mathbf{b}}^{\epsilon_{k_n}}(\tau)\}$, there is a subsequence $\{\hat{\mathbf{b}}^{\epsilon_{h_{k_n}}}(\tau)\}$ of this sequence which converges to $\hat{\mathbf{b}}(\tau)$. Let $\hat{\mathbf{x}}^{\epsilon_n}(\tau) = \mathbf{x}(\tau, \hat{\mathbf{b}}^{\epsilon_n}(\tau), p^{\epsilon_n}(\hat{\mathbf{b}}^{\epsilon_n}))$, for each $n = 1, 2, \dots$, and $\hat{\mathbf{x}}(\tau) = \mathbf{x}(\tau, \hat{\mathbf{b}}(\tau), \hat{p})$. Then, the subsequence $\{\hat{\mathbf{x}}^{\epsilon_{h_{k_n}}}(\tau)\}$ of the sequence $\{\hat{\mathbf{x}}^{\epsilon_n}(\tau)\}$ converges to $\hat{\mathbf{x}}(\tau)$ as the sequence $\{\hat{\mathbf{b}}^{\epsilon_{h_{k_n}}}(\tau)\}$ converges to $\hat{\mathbf{b}}(\tau)$ and the sequence $\{\hat{p}^{\epsilon_{h_{k_n}}}\}$ converges to \hat{p} , with $\hat{p}^{\epsilon_{h_{k_n}}} \gg 0$, for each $n = 1, 2, \dots$, and $\hat{p} \gg 0$. But then, it must be that $\hat{\mathbf{x}}^1(\tau) = 0$ as $\hat{\mathbf{b}}_{21}(\tau) = 0$ and $\hat{\mathbf{x}}(\tau) \in \mathbf{X}^0(\tau, \hat{p})$ as $\hat{\mathbf{x}}^{\epsilon_{h_{k_n}}}(\tau) \in \mathbf{X}^0(\tau, p^{\epsilon_{h_{k_n}}})$, for each $n = 1, 2, \dots$, and the correspondence $\mathbf{X}^0(\tau, \cdot)$ is upper hemicontinuous, by the argument used in Debreu (1982) p. 721. Therefore, we have that $\frac{\partial u_\tau(\hat{\mathbf{x}}(\tau))}{\partial x^1} = +\infty$ as 2 and 1 stand in the relation Q and $\frac{\partial u_\tau(\hat{\mathbf{x}}(\tau))}{\partial x^1} \leq \lambda \hat{p}^1$, by the necessary conditions of the Kuhn-Tucker theorem. Moreover, it must be that $\hat{\mathbf{x}}^2(\tau) = \mathbf{w}^2(\tau) > 0$ as $u_\tau(\cdot)$ is strongly monotone, by Assumption 2, and $\hat{p} \mathbf{w}(\tau) > 0$. Then, $\frac{\partial u_\tau(\hat{\mathbf{x}}(\tau))}{\partial x^2} = \lambda \hat{p}^2$, by the necessary conditions of the Kuhn-Tucker theorem. But then, $\frac{\partial u_\tau(\hat{\mathbf{x}}(\tau))}{\partial x^2} = +\infty$ as $\lambda = +\infty$, contradicting the assumption that $u_\tau(\cdot)$ is continuously differentiable. Therefore, we can conclude that $\tilde{\mathbf{b}}_{21} > 0$. Then, we must also have that $\tilde{\mathbf{b}}_{12} > 0$ as $\tilde{\mathbf{B}}$ is completely reducible. But then, $\tilde{\mathbf{B}}$ is irreducible. Consider a trader $\tau \in T_1$. The matrix $\tilde{\mathbf{B}} \setminus b(\tau)$ is irreducible as $\tilde{\mathbf{b}}_{21} \setminus b(\tau) > 0$, by the previous argument. Consider a trader $\tau \in T_0$. The matrix $\tilde{\mathbf{B}} \setminus b(\tau)$ is irreducible as $\tilde{\mathbf{B}} = \tilde{\mathbf{B}} \setminus b(\tau)$. Then, the matrix $\tilde{\mathbf{B}} \setminus b(\tau)$ is irreducible, for each $t \in T$. But then, from the same argument used by Busetto et al. (2011) in their existence proof (Cases 1 and 3), it follows that $u_t(\mathbf{x}(t, \hat{\mathbf{b}}(t), p(\hat{\mathbf{b}}))) \geq u_t(\mathbf{x}(t, b(t), p(\hat{\mathbf{b}} \setminus b(t))))$, for each

$b(t) \in \mathbf{B}(t)$ and for each $t \in T$. Hence, $\hat{\mathbf{b}}$ is a Cournot-Nash equilibrium. ■

4 Optimality

We now consider the Pareto optimality of a Cournot-Nash allocation. Shapley and Shubik (1977) first raised the question of Pareto optimality of Cournot-Nash allocations in the prototypical strategic market games they introduced. Nevertheless, their analysis was mainly based on examples drawn in an Edgeworth box. Then, some more theoretical results about the Pareto optimality of Cournot-Nash allocations of strategic market games were obtained, both for exchange economies with a finite number of traders and with an atomless continuum of traders, by Dubey (1980), Dubey et al. (1980), Aghion (1985), Dubey and Rogawski (1990), among others. These theorems were obtained in a framework of differential topology and hold generically.⁶ Here, we extend the analysis of Pareto optimality to our mixed model and we obtain a general result which characterizes Pareto optimal Cournot-Nash allocations as Walras allocations. To this end, we need the following further definitions. An allocation \mathbf{x} is said to be individually rational if $u_t(\mathbf{x}(t)) \geq u_t(\mathbf{w}(t))$, for each $t \in T$. An allocation \mathbf{x} is said to be Pareto optimal if there is no allocation \mathbf{y} such that $u_t(\mathbf{y}(t)) > u_t(\mathbf{x}(t))$, for each $t \in T$. An efficiency equilibrium is a pair (p, \mathbf{x}) , consisting of a price vector p and an allocation \mathbf{x} , such that $u_t(\mathbf{x}(t)) \geq u_t(y)$, for all $y \in \{x \in R_+^2 : px = p\mathbf{x}(t)\}$, for each $t \in T$. Moreover, we need to introduce the following assumption.

Assumption 4'. *There is a coalition \bar{T} , with $\bar{T} \subset T_0$, such that $u_t(\cdot)$ is differentiable, additively separable, and $\frac{dv_t^j(0)}{dx^j} = +\infty$, $j = 1, 2$, for each $t \in \bar{T}$.*

It is straightforward to verify that Assumption 4' is stricter than Assumption 4 as Assumption 4' implies Assumption 4 but the converse does not hold. This restriction is needed to guarantee that, at a Pareto optimal Cournot-Nash allocation, Cournot-Nash equilibrium prices are equal, up to a scalar multiple, to efficiency equilibrium prices. We can now state and prove our optimality theorem which establishes an equivalence between the set of Pareto optimal Cournot-Nash allocations and the set of Cournot-Nash allocations which also are Walras allocations.

⁶For a discussion of this literature, see Giraud (2003), p. 359 and p. 365).

Theorem 2. Under Assumptions 1, 2, 3, and 4', let $\hat{\mathbf{b}}$ be a Cournot-Nash equilibrium and let $\hat{p} = p(\hat{\mathbf{b}})$ and $\hat{\mathbf{x}}(t) = \mathbf{x}(t, \hat{\mathbf{b}}(t), p(\hat{\mathbf{b}}))$, for each $t \in T$. Then, $\hat{\mathbf{x}}$ is Pareto optimal if and only if the pair $(\hat{p}, \hat{\mathbf{x}})$ is a Walras equilibrium.

Proof. Let $\hat{\mathbf{b}}$ be a Cournot-Nash equilibrium and let $\hat{p} = p(\hat{\mathbf{b}})$ and $\hat{\mathbf{x}}(t) = \mathbf{x}(t, \hat{\mathbf{b}}(t), p(\hat{\mathbf{b}}))$, for each $t \in T$. Suppose that $\hat{\mathbf{x}}$ is Pareto optimal. We adapt to our framework the argument used by Shitovitz (1973) to prove the corollary to his Lemma 2. It is straightforward to verify that $\hat{\mathbf{x}}$ is individually rational. Let $\hat{\mathbf{G}} \rightarrow \mathcal{P}(R^2)$ be a correspondence such that $\hat{\mathbf{G}}(t) = \{x - \hat{\mathbf{x}}(t) : x \in R_+^2 \text{ and } u_t(x) > u_t(\hat{\mathbf{x}}(t))\}$, for each $t \in T$. Moreover, let $\int_T \hat{\mathbf{G}}(t) d\mu = \{\int_T \hat{\mathbf{g}}(t) d\mu : \hat{\mathbf{g}}(t) \text{ is integrable and } \hat{\mathbf{g}}(t) \in \hat{\mathbf{G}}(t), \text{ for each } t \in T\}$. The set $\{x \in R_+^2 : u_t(x) \geq u_t(\hat{\mathbf{x}})\}$ is convex as $u_t(\cdot)$ is quasi-concave, by Assumption 2, for each $t \in T_1$. Then, it is straightforward to verify that the set $\hat{\mathbf{G}}(t)$ is convex, for each $t \in T_1$. But then, $\int_T \hat{\mathbf{G}}(t) d\mu$ is convex, by Theorem 1 in Shitovitz (1973). We now prove that $0 \notin \int_T \hat{\mathbf{G}}(t) d\mu$. Suppose that $0 \in \int_T \hat{\mathbf{G}}(t) d\mu$. Then, there is an assignment \mathbf{y} such that $u_t(\mathbf{y}(t)) > u_t(\hat{\mathbf{x}}(t))$, for each $t \in T$, which is an allocation as $\int_T \mathbf{y}(t) d\mu = \int_T \hat{\mathbf{x}}(t) d\mu = \int_T \mathbf{w}(t) d\mu$. But then, $\hat{\mathbf{x}}$ is not Pareto optimal, a contradiction. Therefore, it must be that $0 \notin \int_T \hat{\mathbf{G}}(t) d\mu$. Then, there exists a vector \tilde{p} such that $\tilde{p} \in R^2$, ($\tilde{p} \neq 0$), and $\tilde{p} \int_T \hat{\mathbf{G}}(t) d\mu \geq 0$, by the supporting hyperplane theorem. But then, the pair $(\tilde{p}, \hat{\mathbf{x}})$ is an efficiency equilibrium, by Lemma 2 in Shitovitz (1973). We have that $\hat{\mathbf{x}}(t) \in \mathbf{X}^0(t, \tilde{p})$, by the same argument used by Codognato and Ghosal (2000) to prove their Theorem 2, for each $t \in T_0$. Consider a trader $\tau \in \bar{T}$ and suppose that either $\hat{\mathbf{x}}^1(t) = 0$ or $\hat{\mathbf{x}}^2(t) = 0$. Then, the necessary Kuhn-Tucker conditions lead, *mutatis mutandis*, to the same contradiction as in the proof of Theorem 1, by Assumption 4'. But then, we have that $\hat{\mathbf{x}}(t) \gg 0$. Therefore, it must be that

$$\frac{\partial u_t(\hat{\mathbf{x}}(t))}{\partial x^1} = \frac{\hat{p}^1}{\hat{p}^2},$$

for each $t \in \bar{T}$. It must also be that

$$\frac{\partial u_t(\hat{\mathbf{x}}(t))}{\partial x^1} = \frac{\tilde{p}^1}{\tilde{p}^2},$$

as the pair $(\tilde{p}, \hat{\mathbf{x}})$ is an efficiency equilibrium, for each $t \in \bar{T}$. Then, there exists a real number $\theta > 0$ such that $\hat{p}^1 = \theta \tilde{p}^1$ and $\hat{p}^2 = \theta \tilde{p}^2$. But then, $\hat{\mathbf{x}}$ is

such that $\hat{p}\hat{\mathbf{x}}(t) = \hat{p}\mathbf{w}(t)$ and $u_t(\hat{\mathbf{x}}(t)) \geq u_t(y)$, for all $y \in \{x \in R_+^2 : \hat{p}x = \hat{p}\mathbf{w}(t)\}$, for each $t \in T$. Therefore, the pair $(\hat{p}, \hat{\mathbf{x}})$ is a Walras equilibrium. Suppose now that the pair $(\hat{p}, \hat{\mathbf{x}})$ is a Walras equilibrium. Then, $\hat{\mathbf{x}}$ is Pareto optimal, by the first fundamental theorem of welfare economics. Hence, $\hat{\mathbf{x}}$ is Pareto optimal if and only if the pair $(\hat{p}, \hat{\mathbf{x}})$ is a Walras equilibrium. ■

We now consider the relationship between the set of Cournot-Nash allocations, the core, and the set of Walras allocations. We say that an allocation \mathbf{y} dominates an allocation \mathbf{x} via a coalition S if $u_t(\mathbf{y}(t)) > u_t(\mathbf{x}(t))$, for each $t \in S$, and $\int_S \mathbf{y}(t) d\mu = \int_S \mathbf{w}(t) d\mu$. The core is the set of all allocations which are not dominated via any coalition. The following corollary is a straightforward consequence of Theorem 2.

Corollary 1. *Under Assumptions 1, 2, 3, and 4', let $\hat{\mathbf{b}}$ be a Cournot-Nash equilibrium and let $\hat{p} = p(\hat{\mathbf{b}})$ and $\hat{\mathbf{x}}(t) = \mathbf{x}(t, \hat{\mathbf{b}}(t), p(\hat{\mathbf{b}}))$, for each $t \in T$. Then, $\hat{\mathbf{x}}$ is in the core if and only if the pair $(\hat{p}, \hat{\mathbf{x}})$ is a Walras equilibrium.*

Proof. Let $\hat{\mathbf{b}}$ be a Cournot-Nash equilibrium and let $\hat{p} = p(\hat{\mathbf{b}})$ and $\hat{\mathbf{x}}(t) = \mathbf{x}(t, \hat{\mathbf{b}}(t), p(\hat{\mathbf{b}}))$, for each $t \in T$. Suppose that $\hat{\mathbf{x}}$ is in the core. Then, $\hat{\mathbf{x}}$ is Pareto optimal. But then, the pair $(\hat{p}, \hat{\mathbf{x}})$ is a Walras equilibrium, by Theorem 2. Suppose that the pair $(\hat{p}, \hat{\mathbf{x}})$ is a Walras equilibrium. Then, $\hat{\mathbf{x}}$ is in the core, by the same argument used by Aumann (1964) in the proof of his main theorem. Hence, $\hat{\mathbf{x}}$ is in the core if and only if the pair $(\hat{p}, \hat{\mathbf{x}})$ is a Walras equilibrium. ■

Codognato et al. (2015) provided a necessary and sufficient condition for a Cournot-Nash allocation to be a Walras allocation. This characterization result requires a further assumption.

Assumption 5. $u_t : R_+^2 \rightarrow R$ is differentiable, for each $t \in T_1$.

The following proposition provides a characterization of Pareto optimal Cournot-Nash allocations.

Proposition 2. *Under Assumptions 1, 2, 3, 4', and 5, let $\hat{\mathbf{b}}$ be a Cournot-Nash equilibrium and let $\hat{p} = p(\hat{\mathbf{b}})$ and $\hat{\mathbf{x}}(t) = \mathbf{x}(t, \hat{\mathbf{b}}(t), p(\hat{\mathbf{b}}))$, for each $t \in T$. Then, $\hat{\mathbf{x}}$ is Pareto optimal if and only if $\hat{\mathbf{x}}^1(t) = 0$ or $\hat{\mathbf{x}}^2(t) = 0$, for each $t \in T_1$.*

Proof. Let $\hat{\mathbf{b}}$ be a Cournot-Nash equilibrium and let $\hat{p} = p(\hat{\mathbf{b}})$ and $\hat{\mathbf{x}}(t) = \mathbf{x}(t, \hat{\mathbf{b}}(t), p(\hat{\mathbf{b}}))$, for each $t \in T$. Suppose that $\hat{\mathbf{x}}$ is Pareto optimal. Then, the pair $(\hat{p}, \hat{\mathbf{x}})$ is a Walras equilibrium, by Theorem 2. But then, $\hat{\mathbf{x}}^1(t) = 0$ or $\hat{\mathbf{x}}^2(t) = 0$, for each $t \in T_1$, by Theorem 4 in Codognato et al. (2015).

Suppose that $\hat{\mathbf{x}}^1(t) = 0$ or $\hat{\mathbf{x}}^2(t) = 0$, for each $t \in T_1$. Then, the pair $(\hat{p}, \hat{\mathbf{x}})$ is a Walras equilibrium, by Theorem 4 in Codognato et al. (2015). But then, $\hat{\mathbf{x}}$ is Pareto optimal, by Theorem 2. Hence, $\hat{\mathbf{x}}$ is Pareto optimal if and only if $\hat{\mathbf{x}}^1(t) = 0$ or $\hat{\mathbf{x}}^2(t) = 0$, for each $t \in T_1$. ■

The following corollary provides a characterization of Cournot-Nash allocations which are in the core.

Corollary 2. *Under Assumptions 1, 2, 3, 4', and 5, let $\hat{\mathbf{b}}$ be a Cournot-Nash equilibrium and let $\hat{p} = p(\hat{\mathbf{b}})$ and $\hat{\mathbf{x}}(t) = \mathbf{x}(t, \hat{\mathbf{b}}(t), p(\hat{\mathbf{b}}))$, for each $t \in T$. Then, $\hat{\mathbf{x}}$ is the core if and only if $\hat{\mathbf{x}}^1(t) = 0$ or $\hat{\mathbf{x}}^2(t) = 0$, for each $t \in T_1$.*

Proof. Let $\hat{\mathbf{b}}$ be a Cournot-Nash equilibrium and let $\hat{p} = p(\hat{\mathbf{b}})$ and $\hat{\mathbf{x}}(t) = \mathbf{x}(t, \hat{\mathbf{b}}(t), p(\hat{\mathbf{b}}))$, for each $t \in T$. Suppose that $\hat{\mathbf{x}}$ is in the core. Then, $\hat{\mathbf{x}}$ is Pareto optimal. But then, $\hat{\mathbf{x}}^1(t) = 0$ or $\hat{\mathbf{x}}^2(t) = 0$, for each $t \in T_1$, by Proposition 2. Suppose that $\hat{\mathbf{x}}^1(t) = 0$ or $\hat{\mathbf{x}}^2(t) = 0$, for each $t \in T_1$. Then the pair $(\hat{p}, \hat{\mathbf{x}})$ is a Walras equilibrium, by Theorem 4 in Codognato et al. (2015). But then, $\hat{\mathbf{x}}$ is in the core, by the same argument used by Aumann (1964) in the proof of his main theorem. ■

Examples 6, 7, 8, and 9 in Codognato et al. (2015) show that Theorem 2, Proposition 2, and Corollaries 1 and 2 hold non-vacuously.

5 Discussion of the model

We now discuss some issues related to the existence and optimality of Cournot-Nash equilibria. It is straightforward to show, using Theorem 5 in Codognato et al. (2015), that, in our mixed bilateral oligopoly framework, under Assumptions 1, 2, and 3, the set of Cournot-Nash allocations of the Shapley window model coincides with the set of the Cournot-Nash allocations of the model with commodity money proposed by Dubey and Shubik (1978) and of its generalization to complete markets proposed by Amir et al. (1990). Therefore, all the results obtained in this paper also hold for these models. Let us now further discuss some features of this class of models, in the bilateral oligopoly framework, and the results we have obtained.

Busetto et al. (2017) considered a mixed version of the Shapley window model for exchange economies with l commodities. In their Theorem 2, they proved the existence of a Cournot-Nash equilibrium under their Assumptions 1, 2, 3, and 4. In this paper, we have considered a bilateral oligopoly version of the model analyzed by Busetto et al. (2017). Nevertheless, Theorem 1, our new existence theorem, is not just a two-commodity case of Theorem 2

in Busetto et al. (2017). While Assumptions 2 and 3 are the same in Busetto et al. (2017) and here, Assumptions 1 and 4 differ. We shall now analyze more in detail the difference between the two versions of Assumptions 1 and 4 and the role they play in the existence proofs.

In our bilateral framework, Assumption 1 in Busetto et al. (2017) could be restated as follows.

Assumption 1'. *There is a coalition S such that $\mu(S \cap T) > 0$, $\mu(S^c \cap T) > 0$, $\mathbf{w}^1(t) > 0$, $\mathbf{w}^2(t) = 0$, for each $t \in S$, $\mathbf{w}^1(t) = 0$, $\mathbf{w}^2(t) > 0$, for each $t \in S^c$. Moreover, $\int_{T_0} \mathbf{w}(t) d\mu \gg 0$.*

It is clear that if an initial assignment \mathbf{w} satisfies Assumption 1', then it also satisfies our Assumption 1. However, the cases where $\mu(S \cap T_0) = 0$ or $\mu(S^c \cap T_0) = 0$, that is, where there are only atoms holding one of the two commodities are ruled out by Assumption 1'. Therefore, the price convergence theorem proved by Busetto et al. (2017), their Theorem 1, cannot be used in the proof of our existence theorem as it only holds under Assumption 1'. In order to cover the case where one of the two commodities is held only by atoms, we have proved a price convergence lemma which combines the argument used by Busetto et al. (2017) in their Theorem 1 with that used by Dubey and Shubik (1978) in their Lemma 2, which provides a price convergence result in the purely atomic case. Moreover, we have extended the proof of Lemma 2 in Dubey and Shubik (1978), which holds under the assumption of concave utility functions, to cover the case of quasi-concave utility functions as required by our Assumption 2.

Assumption 4 in Busetto et al. (2017) is based on a relation between commodities, their relation C , which coincides, in our bilateral framework, with our relation Q . It requires that the set of commodities is strongly connected through traders' characteristics and it could be restated, in our framework, as follows.

Assumption 4''. *Commodities 1 and 2 and commodities 2 and 1 stand in relation Q .*

It is clear that Assumption 4'' implies our Assumption 4 but the converse does not hold. Therefore, in order to show the irreducibility of the matrix $\hat{\mathbf{B}}$, in the proof of Theorem 1, bilateral oligopoly allows us to impose condition Q only for one of the two ordered pairs generated by commodities 1 and 2.

Therefore, we can conclude, from the previous discussion, that our existence theorem holds under less restrictive conditions than the existence theorem in Busetto et al. (2017).

So far we have kept the assumption that $\mu(T_1) > 0$ and $\mu(T_0) > 0$. Suppose first that $\mu(T_1) = 0$. Then, it is possible to adapt, to the atomless bilateral framework we obtain, the equivalence result proved by Codognato and Ghosal (2000), to prove the existence of a Cournot-Nash equilibrium in this case.

Proposition 3. *Let $\mu(T_1) = 0$. Under Assumptions 1, 2, 3, and 4, there exists a Cournot-Nash equilibrium $\hat{\mathbf{b}}$.*

Proof. Under Assumptions 1, 2, 3, and 4, there exists a Walras equilibrium $(\hat{p}, \hat{\mathbf{x}})$, by Theorem 9 in Debreu (1982), a generalization of an existence theorem for exchange economies with an atomless continuum of traders first proved by Aumann (1966). Then, there exists a strategy selection $\hat{\mathbf{b}}$ such that $\hat{p} = p(\hat{\mathbf{b}})$ and $\hat{\mathbf{x}}(t) = \mathbf{x}(t, \hat{\mathbf{b}}(t), p(\hat{\mathbf{b}}))$, for each $t \in T$, which is a Cournot-Nash equilibrium, by the argument used by Codognato and Ghosal (2000) to prove their Theorem 2, replacing their Assumption 4, which requires that the set of commodities is a net, with our Assumption 4. ■

Suppose now that $\mu(T_0) = 0$. Then, it is possible to adapt, to the purely atomic bilateral framework we obtain, the existence result proved by Bloch and Ferrer (2001). To this end, let us introduce the following restrictions of Assumptions 2 and 5, respectively.

Assumption 2'. $u_t : R_+^2 \rightarrow R$ is continuous, strongly monotone, and strictly concave, for each $t \in T_1$.

Assumption 5'. $u_t(\cdot)$ is differentiable, additively separable, and $\frac{dv_t^j(0)}{dx^j} = +\infty$, $j = 1, 2$, for each $t \in T_1$.

We can then state the existence result proved by Bloch and Ferrer (2001).

Proposition 4. *Let $\mu(T_0) = 0$. Under Assumptions 1, 2', 3, and 5', there exists a Cournot-Nash equilibrium $\hat{\mathbf{b}}$.*

Proof. We have that $\text{card}(S \cap T_1) \geq 2$ and $\text{card}(S^c \cap T_1) \geq 2$ as $\mu(T_0) = 0$, by Assumption 1. Then, there exists a Cournot-Nash equilibrium $\hat{\mathbf{b}}$, by the same argument used by Bloch and Ferrer (2001) to prove their Proposition 2, which can also be extended to the case where T_1 contains countably infinite atoms by means of the product topology. ■

It could be worth investigating, if Proposition 4 holds under weaker assumptions.⁷

⁷It would also be worth comparing the existence proof provided by Bloch and Ferrer

We have already noticed, in Section 4, that Assumption 4' is more restrictive than Assumption 4. We leave for further research a possible answer to the question whether Theorem 2 holds under less restrictive or alternative assumptions.

Dubey and Rogawski (1990) showed that, for strategic market games with a finite number of traders, Cournot-Nash equilibria are generically not Pareto optimal in utility functions. In future work, we could verify if this result would hold in our bilateral oligopoly framework as it would imply, by Theorem 2, that the “probability” that a Cournot-Nash allocation is a Walras allocation is null.

In Section 2, we have provided a definition of a Cournot-Nash equilibrium referring explicitly to irreducible matrices. This definition applies only to active Cournot-Nash equilibria according to the definition of Sahi and Yao (1989). Nevertheless, in the Shapley window model, as in all other strategic market games, the strategy selection $\hat{\mathbf{b}}$ such that $\hat{\mathbf{b}}(t) = 0$, for each $t \in T$, is a Cournot-Nash equilibrium, usually called trivial equilibrium. This raises the question whether, under Assumptions 1-4, the allocation corresponding to the trivial Cournot-Nash equilibrium, namely the initial assignment \mathbf{w} , may be Pareto optimal. The following proposition provides a negative answer to this question.

Proposition 5. *Under Assumptions 1, 2, 3, and 4, the allocation \mathbf{w} is not Pareto optimal.*

Proof. Suppose that \mathbf{w} is Pareto optimal. Then, there exists a price vector \tilde{p} such that the pair (\tilde{p}, \mathbf{w}) is an efficiency equilibrium, by the same argument used in the proof of Theorem 2. But then, the pair (\tilde{p}, \mathbf{w}) is a Walras equilibrium. Therefore, for commodities which stand in the relation Q , the necessary Kuhn-Tucker conditions lead to the same contradiction as in the proof of Theorem 1. Hence, the allocation \mathbf{w} is not Pareto optimal. ■

6 Conclusion

In this paper, we have shown, in Theorem 1, the existence of a Cournot-Nash equilibrium for the mixed bilateral oligopoly version of the Shapley window model first analyzed by Codognato et al. (2015). Then, we have proved, in Theorem 2, that a Cournot-Nash allocation is Pareto optimal if

(2001) with that given by Dickson and Hartley (2008), based on another approach to bilateral oligopoly.

and only if it is a Walras allocation. The proof of this theorem is based on a corollary in Shitovitz (1973) which shows that the first and second welfare theorem still hold in mixed exchange economies. Codognato et al. (2015) showed, in their main theorem, that, under a further assumption of differentiability of atoms' utility functions, the condition which characterizes the nonempty intersection of the sets of Walras and Cournot-Nash allocations requires that each atom demands a null amount of one commodity. Combining this result with our Theorem 2 we have obtained, as a proposition, a characterization of the optimality of Cournot-Nash equilibria which requires that at a Pareto optimal Cournot-Nash equilibrium each atom demands a null amount of one commodity. Recasting antitrust analysis in the bilateral oligopoly framework, in further research, we could use these results as a first step to analyze competition policy in a general equilibrium framework. As we said, the results we have obtained in this paper also holds for the main prototypes of strategic market games inspired by Shapley and Shubik (1977). In forthcoming research, we would like to verify if they also hold for another class of models introduced by Postlewaite and Schmeidler (1978) and further analyzed by Peck et al. (1992), Koutsougeras and Ziros (2008), Koutsougeras (2009), among others.

References

- [1] Aghion P. (1985), "On the generic inefficiency of differentiable market games," *Journal of Economic Theory* **37**, 126-146.
- [2] Amir R., Bloch F. (2009), "Comparative statics in a simple class of strategic market games," *Games and Economic Behavior* **65**, 7-24.
- [3] Amir R., Sahi S., Shubik M., Yao S. (1990), "A strategic market game with complete markets," *Journal of Economic Theory* **51**, 126-143.
- [4] Artstein Z. (1979), "A note on Fatou's lemma in several dimensions," *Journal of Mathematical Economics* **6**, 277-282.
- [5] Aumann R.J. (1964), "Markets with a continuum of traders," *Econometrica* **32**, 39-50.
- [6] Aumann R.J. (1966), "Existence of competitive equilibria in markets with a continuum of traders," *Econometrica* **24**, 1-17.

- [7] Bloch F., Ghosal S. (1997), "Stable trading structures in bilateral oligopolies," *Journal of Economic Theory* **74**, 368-384.
- [8] Bloch F., Ferrer H. (2001), "Trade fragmentation and coordination in strategic market games," *Journal of Economic Theory* **101**, 301-316.
- [9] Busetto F., Codognato G., Ghosal S. (2011), "Noncooperative oligopoly in markets with a continuum of traders," *Games and Economic Behavior* **72**, 38-45.
- [10] Busetto F., Codognato G., Ghosal S., Julien L., Tonin S. (2017), "Non-cooperative oligopoly in markets with a continuum of traders and a strongly connected set of commodities," *Games and Economic Behavior*, forthcoming.
- [11] Codognato G., Ghosal S. (2000), "Cournot-Nash equilibria in limit exchange economies with complete markets and consistent prices," *Journal of Mathematical Economics* **34**, 39-53.
- [12] Codognato G., Ghosal S., Tonin S. (2015), "Atomic Cournotian traders may be Walrasian," *Journal of Economic Theory* **159**, 1-14.
- [13] Debreu G. (1982), "Existence of competitive equilibrium," in Arrow K.J., Intriligator H.D. (eds), *Handbook of mathematical economics*, Elsevier, Amsterdam.
- [14] Dickson A., Hartley R. (2008), "The strategic Marshallian cross," *Games and Economic Behavior* **64**, 514-532.
- [15] Dubey P. (1980), "Nash equilibria of market games: finiteness and inefficiency," *Journal of Economic Theory* **22**, 363-376.
- [16] Dubey P., Rogawski J.D. (1990), "Inefficiency of smooth market mechanisms," *Journal of Mathematical Economics* **19**, 285-304.
- [17] Dubey P., Shubik M. (1978), "The noncooperative equilibria of a closed trading economy with market supply and bidding strategies," *Journal of Economic Theory* **17**, 1-20.
- [18] Dubey P., Mas-Colell A., Shubik M. (1980), "Efficiency properties of strategic market games: an axiomatic approach," *Journal of Economic Theory* **22**, 339-362.

- [19] Gabszewicz J.J., Michel P. (1997), "Oligopoly equilibrium in exchange economies," in Eaton B.C., Harris R. G. (eds), *Trade, technology and economics. Essays in honour of Richard G. Lipsey*, Edward Elgar, Cheltenham.
- [20] Giraud G. (2003), "Strategic market games: an introduction," *Journal of Mathematical Economics* **39**, 355-375.
- [21] Koutsougeras L.C. (2009), "Convergence of strategic behavior to price taking," *Games and Economic Behavior* **65**, 234-241.
- [22] Koutsougeras L.C., Ziros N. (2008), "A three way equivalence," *Journal of Economic Theory* **139**, 380-391.
- [23] Kreps D. (2012), *Microeconomic foundations I: choice and competitive markets*, Princeton University Press, Princeton.
- [24] Peck J., Shell K., Spear S.E. (1992), "The market game: existence and structure of equilibrium," *Journal of Mathematical Economics* **21**, 271-299.
- [25] Postlewaite A., Schmeidler D. (1978), "Approximate efficiency of non-Walrasian Nash equilibrium," *Econometrica* **46**, 127-135.
- [26] Sahi S., Yao S. (1989), "The noncooperative equilibria of a trading economy with complete markets and consistent prices," *Journal of Mathematical Economics* **18**, 325-346.
- [27] Shapley L.S., Shubik M. (1977), "Trade using one commodity as a means of payment," *Journal of Political Economy* **85**, 937-968.
- [28] Shitovitz B. (1973), "Oligopoly in markets with a continuum of traders," *Econometrica* **41**, 467-501.